

## Mouse Anti Human CD19 Purified

### PRODUCT INFORMATION

<b>CLONE:</b>	HI19a
<b>ISOTYPE:</b>	Mouse IgG1, $\kappa$
<b>WS.No.:</b>	V CD19.11
<b>CATALOG#:</b>	A6361/A6361-bulk
<b>CONTENTS:</b>	Purified antibody buffered in 10mM PBS (pH 7.0) with 0.1% NaN <sub>3</sub> .

### DESCRIPTION

CD19 McAb recognizes a 95 KD type I transmembrane glycoprotein which is restricted B cell antigen. CD19 antigen is expressed on normal and neoplastic B cells and also in some the bone marrow cells. CD19 expression by B progenitor cells is presumably at late pro-B or early pre-B stages around the time of Ig heavy chain rearrangement. Expression persists during all stages of B cell maturation and is lost on terminal differentiation to plasma cells. CD19 antigen is also found on the follicular dendritic cells and the early cells of myelomonocytic lineage but not on normal T cells, NK cells, monocytes, granulocytes, erythrocytes and platelets. In normal peripheral blood, 8-20% of lymphocytes express CD19 antigen. CD19 antigen plays a role in regulating B cell proliferation.

### PREPARATION

The monoclonal antibody is purified from ascites by protein G affinity chromatography.

### USAGE

The purified reagent is effective for indirect immunofluorescence staining of human cells for flow cytometric analysis and is tested for immunohistochemical staining of acetone-fixed frozen sections.

### STORAGE

For purified forms, long term storage at -20°C. Avoid freeze and thaw cycles.

### REFERENCES

Schlossman, S., L.Bloumsell, W.Gilks, et al., eds. 1995. Leucocyte Typing V: White Cell Differentiation Antigens. P:491、507、2004 Oxford University Press, New York.

Use For Research Only